

INFECTIOUS GASTROENTERITIS TESTING GUIDE

| Clinical History | Routine Faecal Culture | Giardia & Cryptos- poridium antigen | Full Extended Parasite examination (2 samples) | Clostridium difficile toxin testing | Viral testing for gastro | Other points Note relevant details on 'clinical details' to help the lab guide the testing |
|---|------------------------------|--|---|--|--------------------------------|---|
| Acute diarrhoea only | No testing | | | | | Fluid electrolyte replacement only, dependent on age and other factors |
| Bloody diarrhoea or febrile | ✓ | | | | | One sample - 85% sensitive when bacterial cause |
| Chronic diarrhoea | ✓ | ✓ | ✓ | | | Note if overseas travel and where if one main location |
| Children ≤ 5 years | ✓ | | | | * | * Consider rotavirus also especially if outbreak and/or childcare attendance |
| Post antibiotic &/or hospital &/or LTCF | | | | ✓ | | If positive toxin, do not retest |
| > 65 years | ✓ | | | * | | * Consider <i>C difficile</i> also if Abs, debilitation, or has been in healthcare facility |
| Overseas travel or immigration | ✓ | ✓ | ✓ | | | , |
| Immunocompro- mised | ✓ | ✓ | ✓ | ✓ | | |
| Rural or animal exposure | ✓ | ✓ | | | | |
| Child care attendance | ✓ | ✓ | | | * | * Consider viral testing especially if > 1 case |
| Healthcare worker | ✓ | | | ✓ | ✓ | Extent of testing guided by patient contact type |
| Food handler | ✓ | | | | * | Plus consider viral if > 1 case |
| Raw seafood | ✓ | | | | ✓ | Note details on form, extra bacterial testing will be done e.g. <i>Vibrio spp</i> |
| Outbreak | ✓ | | | * | * | * Consider extra testing dependent on point source (food poisoning), healthcare association (C. diff), or institutional (viral) |