1 START THIS CHART IF THE PATIENT LOOKS UNWELL OR HAS ABNORMAL PHYSIOLOGY

RISK FACTORS FOR SEPSIS INCLUDE:

Age > 73
Impaired immunity (e.g. diabetes, steroids, chemotherapy)

Recent trauma / surgery / invasive procedure Indwelling lines / IVDU / broken skin

02 COULD THIS BE DUE TO AN INFECTION?

LIKELY SOURCE:

 \square $\Delta \sigma \rho > 75$

Respiratory
Brain

Urine
Surgical

Skin / joint / wound
Other

YES

☐ Indwelling device

NO

SEPSIS UNLIKELY, CONSIDER OTHER DIAGNOSIS

13 ANY RED FLAG PRESENT?

 \Box Objective evidence of new or altered mental state \Box Systolic BP ≤ 90 mmHg (or drop of >40 from normal)

Heart rate ≥ 130 per minute

Respiratory rate ≥ 25 per minute

Needs O_2 to keep $SpO_2 \ge 92\%$ (88% in COPD)

☐ Non-blanching rash / mottled / ashen / cyanotic

Recent chemotherapy

Not passed urine in 18 hours (<0.5ml/kg/hr if catheterised)

RED FLAG SEPSIS START GP BUNDLE

1 ANY AMBER FLAG PRESENT?

IF UNDER 17 & IMMUNITY IMPAIRED TREAT AS RED FLAG SEPSIS

Relatives concerned about mental status
 Acute deterioration in functional ability
 Immunosuppressed

Trauma / surgery / procedure in last 8 weeks

Respiratory rate 21-24

Systolic BP 91-100 mmHg

Heart rate 91-130 or new dysrhythmia

Temperature <36°C</p>

Clinical signs of wound infection

USE CLINICAL JUDGEMENT TO DETERMINE WHETHER PATIENT CAN BE MANAGED IN COMMUNITY SETTING. IF TREATING IN THE COMMUNITY CONSIDER:

- PLANNED SECOND

ASSESSMENT +/- BLOODS

- SPECIFIC SAFETY NETTING ADVICE

NO AMBER FLAGS : ROUTINE CARE AND GIVE SAFETY-NETTING ADVICE:

CALL 111 IF CONDITION CHANGES OR DETERIORATES.
SIGNPOST TO AVAILABLE RESOURCES AS APPROPRIATE

CALL 999 IF ANY OF: Slurred speech or confusion
Extreme shivering or muscle pain
Passing no urine (in a day)
Severe breathlessness
'I feel I might die'
Skin mottled, ashen, blue or very pale

GP RED FLAG BUNDLE:

THIS IS TIME-CRITICAL - IMMEDIATE ACTION REQUIRED: DIAL 999

AND ARRANGE BLUE LIGHT TRANSFER

COMMUNICATION: Ensure communication of 'Red Flag Sepsis' to crew. Advise crew to pre-alert as 'Red Flag Sepsis'. Where possible a written handover is recommended including observations and antibiotic allergies.



UKST 2020 3.2 PAGE 1 0F 1